


Norovirus

Schools Guide

for early years professionals

This pack contains

- Information on the Norovirus aka, the 'winter vomiting bug'
- Advice on limiting the spread of Norovirus in your school
 - Guidance on how symptoms can be treated
 - A printout to give to parents and carers



Diarrhoea and vomiting in children

There's no specific cure for stomach bugs such as **Norovirus**. See tips to reduce the spread and treat symptoms

<p>Keep kids home from school for</p> <p>48 HOURS</p> <p>after symptoms have passed. This is enough time to let the virus run its course</p> <p>If your child already has a serious illness, or symptoms last longer than a few days, contact your GP to seek advice Visit nhs.uk/norovirus</p> <p><small>NHS choices Food Standards Agency</small></p>	<p>Washing hands frequently and thoroughly helps limit the spread of the virus</p>	<p>25 different strains of Norovirus are known to affect humans. They're the most common cause of stomach bugs in the UK</p>
<p>Home Treatment</p> <p>Paracetamol is useful for fever or aches and pains</p> <p>Give kids plenty of water to stay hydrated and replace lost fluids</p> <p>If your child feels like eating, give them bland foods</p>		

Norovirus schools guide

Norovirus, also known as the winter vomiting bug, is the most common stomach bug in the UK. NHS Choices, in collaboration with the Food Standards Agency, has put together this useful guide and printout to help schools and parents understand Norovirus, from detection to prevention.

What is Norovirus?

Norovirus is highly contagious. It's particularly prevalent in schools and nurseries as the virus can survive for several days on surfaces or objects. But it can affect people of all ages and cause vomiting and diarrhoea.

There is no cure for Norovirus, so it has to be left to run its course. The symptoms are unpleasant and can initially be quite distressing, but they shouldn't last for more than a couple of days. Adults and children with Norovirus symptoms should avoid visiting GP surgeries or hospitals to prevent the further spread of the virus.

Because the virus is highly contagious, **children who have Norovirus symptoms must remain off school or nursery for 48 hours after the last episode of vomiting or diarrhoea to stop the spread of the infection to other children and staff.**

Different types of Norovirus

Norovirus is the most common cause of stomach bugs in the UK, with at least 25 different strains known to affect humans. Each year, it's estimated between 600,000 and 1 million people in the UK catch Norovirus – because there are so many strains, we don't develop immunity to it.

The virus is sometimes called the "winter vomiting bug" as it's more common in winter. However, you can catch the virus at any time of the year.

How do you treat Norovirus?

To treat the symptoms of Norovirus, we recommend the following for children and young people:

- Ensure they drink plenty of water to avoid dehydration.
- Give them liquid paracetamol for any fever or aches and pains.
- If they feel like eating, give them easy to digest foods such as bananas or brown rice.
- Stay at home – there is nothing the GP can prescribe for sickness and diarrhoea, although your local chemist may be able to provide rehydration solutions.
- Contact your GP or NHS 111 to seek advice if the symptoms last longer than a few days, or visit [nhs.uk/norovirus](https://www.nhs.uk/norovirus) for more information.

Preventing the spread of Norovirus

Norovirus is easily spread – you can catch it simply by touching contaminated surfaces or objects.

Early years staff, teachers, school staff, and parents can help prevent the spread of Norovirus by:

- washing hands frequently with household soap, and encouraging children to do the same – especially after going to the toilet
- ensuring any infected child is not sharing things such as toys, blankets, and so on
- keeping any child with Norovirus symptoms home from school and away from other children where possible – children with Norovirus should also avoid contact with vulnerable adults, such as elderly relatives

It is also important that schools and nurseries thoroughly disinfect all surfaces an infected person may have come into contact with whenever possible. Alcohol-based hand gels are not effective against Norovirus.

For further information, visit nhs.uk/norovirus or contact NHS 111.

You may wish to talk to your school nurse about infection control and materials that can be used in schools to increase pupil and parent awareness.

Project background

NHS Choices and the Food Standards Agency have this year conducted a significant research project looking at how we can best tackle some of the issues of misinformation surrounding winter vomiting bugs.

The information in this pack has been produced to empower teachers and school nurses to help stop the spread of Norovirus, and enable parents and guardians to react to the virus effectively and without putting others at risk.

By distributing the information within this pack, you are helping to reduce the spread of a highly contagious virus that puts many NHS services under great pressure and can seriously affect the health of some vulnerable groups. This material forms part of a wider digital campaign to help the public understand Norovirus, to be launched later this winter.

Notes for teachers and school nurses

- The black and white version of the poster has been produced to keep the costs of printing at a minimum should you choose, while we recommend schools add the colour version to their website and signpost parents and guardians to the documents.
- Our user research suggests parents are a key group to target with Norovirus information as they are among the most likely to take their child to a GP or hospital with symptoms of the virus – a major cause of infectious spread.
- General understanding of the term “Norovirus” is low, and it may be more helpful to refer to the symptoms (diarrhoea and vomiting) in communications with parents.

Diarrhoea and vomiting in children

There's no specific cure for stomach bugs such as **Norovirus**. See tips to reduce the spread and treat symptoms

Keep kids home from school for



after symptoms have passed.
This is enough time to let the
virus run its course

If your child already has a serious
illness, or symptoms last longer
than a few days, contact your GP
to seek advice

Visit [nhs.uk/norovirus](https://www.nhs.uk/norovirus)



25

different
strains of
Norovirus
are known to
affect humans.

They're the most common cause
of stomach bugs in the UK

Home Treatment



Paracetamol is
useful for fever or
aches and pains



Give kids plenty of water
to stay hydrated and
replace lost fluids



If your child feels
like eating, give
them bland foods

Diarrhoea and vomiting in children

There's no specific cure for stomach bugs such as **Norovirus**. See tips to reduce the spread and treat symptoms

Keep kids home from school for



after symptoms have passed.
This is enough time to let the
virus run its course

If your child already has a serious
illness, or symptoms last longer
than a few days, contact your GP
to seek advice

Visit [nhs.uk/norovirus](https://www.nhs.uk/norovirus)



25

different
strains of
Norovirus
are known to
affect humans.

They're the most common cause
of stomach bugs in the UK


Home Treatment



Paracetamol is
useful for fever or
aches and pains



Give kids plenty of water
to stay hydrated and
replace lost fluids



If your child feels
like eating, give
them bland foods